





Time period: July 15, 2012 - November 16, 2013

Patient totals: Stringent*: 974; Stage 4: 94






Key:					
Report	Measure Name	Cohort	Target	Cohort Definition	Formula
1	Control Blood Pressure	All patients who meet stringent CKD criteria	130/80	Percentage of patients with average of last 3 BP <= 130/80	Mean of last three systolic and diastolic BP; will be based on last one or two if fewer than three available
2	Control LDL	All patients who meet stringent CKD criteria	<100	Percentage of patients with average of last two LDL measures < 100	Mean of last two LDL; last LDL if only one is available
3	Use ACE/ARB	All patients who meet stringent CKD criteria	Yes	Percentage of patients with a current ACE/ARB prescription	Documentation in EHR/pharmacy of prescription; yes/no for each time period
4	Control HbA1C	Patients who meet stringent CKD criteria and have a Dx code of 250*	<7.0	Percentage of diabetic patients with A1c < 7.0	Last HbA1c;
5	Have Eliminated NSAID/Cox-2 use	All patients who meet stringent CKD criteria	No	Percentage of patients with no current prescription for NSAID/Cox 2	Yes/no for each time period
6	Refer to Nephrologist	Patients with at least one GFR < 30	Yes	Percentage of patients with at least one eGFR < 30 who have been referred to nephrology	Referral documented, if applicable
7	Eliminate Smoking	Patients with smoking status documented	Non / Former smokers	Percentage of patients with a documented smoking status who are non smokers or former smokers	Yes/no for each time period
8	Diagnosis of CKD	Pts who meet stringent CKD criteria	Appropriate CKD Dx	Percentage of patients who meet CKD criteria who have CKD Diagnosis	Dx Codes
9	Annual ACR Measure	Pts who have ICD9 of 250.xx	Annual measurement	Percentage of patients who have MicroCR in prior year	Lab done
10	Annual LDL	Pts who meet stringent CKD criteria	Annual measurement	Percentage of patients who have LDL in prior year	Lab done
11	Annual A1c	Patients with diabetes	Annual measurement	Percentage of patients who have A1c in prior year	Lab done
12	Follow up creatinine measure	Pts who have 1 eGFR < 60	Appropriate Follow-Up Measurement	Percentage of patients who have a follow up creatinine measure > 90 and < 360 days after the first eGFR < 60	Lab done
13	Follow up ACR measure	Pts who have 1 ACR > 30	Appropriate Follow-Up Measurement	Percentage of patients who have a follow up ACR measure > 90 days and < 360 days after the first ACR > 30	Lab done

Notes: stringent CKD criteria = 2 eGFR values >= 15 and/or < 60 at least 90 days apart, with no more than one eGFR < 15






Time period: July 15, 2012 - November 16, 2013

Patient totals: Stringent*: 974; Stage 4: 94






Time period: July 15, 2012 - November 16, 2013

Patient totals: Stringent*: 974; Stage 4: 94



Time period: July 15, 2012 - November 16, 2013

Patient totals: Stringent*: 974; Stage 4: 94



Time period: July 15, 2012 - November 16, 2013

Patient totals: Stringent*: 974; Stage 4: 94

